## ADDENDUM AND CORRIGENDA TO "CLASSIFICATION OF THE SUPERFAMILY SESIOIDEA"

### JOHN B. HEPPNER AND W. D. DUCKWORTH

Department of Entomology, Smithsonian Institution, Washington, D.C.

**ABSTRACT.** The subgenera Dipchasphecia, Ductispina, Paradipsosphecia, Pseudosphecia, and Synansphecia are synonymized with Bembecia in Sesiidae. The names Bembecia reisseri Capuse, Chamaesphecia rangnovi Capuse, Homogyna porphyractis Meyrick, Microsynanthedon tanala Minet, and Sesia okinawana (Matsumura) are added. A number of other changes and corrections are noted.

In a recently completed paper on Sesioidea (Heppner & Duckworth, 1981), a major reference to Sesiidae (Capuse, 1973) was inadvertantly left out of the world review for the family Sesiidae. Thus, we wish to make note of the following taxa as an addition to our review of the superfamily.

The following generic names, proposed as subgenera, are hereby placed as synonyms of *Bembecia*:

# Pseudosphecia Capuse, 1973b:147, new synonymy

Type-species: *Dipsosphecia tenebrosa* Püngeler, 1915, by original designation [described as a subgenus of *Bembecia*].

- Ductispina Capuse, 1973b:149, **new synonymy** Type-species: Dipsosphecia turcmena Bartel, 1912, by original designation [described as a subgenus of Bembecia].
- Paradipsosphecia Capuse, 1973b:150, **new synonymy** Type-species: Dipsosphecia barbara Bartel, 1912, by original designation [described as a subgenus of *Bembecia*].
- Dipchasphecia Capuse, 1973b:161, **new synonymy** Type-species: Dipsosphecia roseiventris Bartel, 1912, by original designation [described as a subgenus of Chamaesphecia; typespecies now placed in Bembecia].

Synansphecia Capuse, 1973b:166, **new synonymy** Type-species: Sesia triannuliformis Freyer, 1845, by original designation [described as a subgenus of Chamaesphecia; type-species now placed in Bembecia].

The following additions and corrections are also noted:

- p. 9. Change setae D2 to D1 and D1 to D2 at the end of the larval diagnosis.
- p. 15. Change Phycodes eucallynta Meyrick, 1937, to Meyrick, 1937a.
- p. 16. Under current taxa, change 1063 to 1066 species and change 83 to 84 species.
- p. 20. Under current taxa, change 782 to 786 species. Under Bembecia, change 58 to 59.
- p. 21. Under Homogyna, change 9 to 10; under Microsynanthedon, change 1 to 2.
- p. 27. Under Sesia, add Spherodoptera [sic] Matsumura, 1931a:1017, [misspelling].

- p. 28. Add Sesia okinawana (Matsumura, 1931a:1017), Spherodoptera [sic], [new combination], as a valid species from the Oriental region. Change Sesia montelli (Löfquist, 1922:82), Aegeria from a valid species to a junior synonym of Sesia bembeciformis (Hübner).
- p. 30. Change Synanthedon danieli Capuse, 1973, to Capuse, 1973a.
- p. 35. Under Chamaesphecia balcanica Zukowsky, 1929, add C. rangnovi Capuse, 1973b:167 [nomen nudum], as a synonym. Change C. corsica (Staudinger, 1856) to a synonym of C. meriaeformis (Boisduval, 1840), since the latter is not a nomen nudum, due to a short description in a footnote. Chamaesphecia meriaeformis becomes the valid name for the species.
- p. 36. Change Chamaesphecia monspeliensis (Staudinger, 1856) to synonymy with C. tengyraeformis (Boisduval, 1840), since the latter is not a nomen nudum, due to a short description in a footnote. Chamaesphecia tengyraeformis becomes the valid name for this species.
- p. 37. Change Scalarignathia Capuse, 1973, and Scalarignathia kaszabi Capuse, 1973, to Capuse, 1973a.
- p. 39. Add *Bembecia reisseri* (Capuse, 1973b:156), *Pyropteron* [new combination], as a valid species from the Palearctic region.

Under B. chrysidiformis (Esper, 1782), B. polistiformis (Boisduval, 1840) is not a nomen nudum due to a short description as a footnote in the original article.

Change Bembecia bestianaeli Capuse, 1973, Bembecia dancaudani Capuse, 1973, Bembecia hannemanni Capuse, 1973, and Bembecia ili Capuse, 1973, in each case to "(Capuse, 1973a), Dipsosphecia [new combination]."

- p. 40. Since Chamaesphecia tengyraeformis (Boisduval) 1840 (Sesia) is now a valid name and not a nomen nudum, Bembecia tengyraeformis (Herrich-Schäffer, 1851) (Sesia) becomes a junior homonym, to be replaced by Bembecia sanguinolenta (Lederer, 1853). The latter name becomes the valid name for this species.
- p. 42. Add Microsynanthedon tanala Minet, 1976:40, for the Ethiopian region.
- p. 43. Add *Homogyna porphyractis* Meyrick, 1937b:119, as a valid species from the Ethiopian region.
- p. 49. Change Brenthia dendronympha Meyrick, 1937, to Meyrick, 1937a.
- p. 50. Change Brenthia spintheristis [sic] Meyrick to B. spintheritis.
- p. 51. Under Millieria dolosana (Herrich-Schäffer) add M. dolosalis (Heydenreich, 1851:63), Choreutis [nomen nudum], as another synonym.
- p. 52. Change Tebenna chrysotacta [sic] (Meyrick) to T. chrysostacta.
- p. 62. Add the Capuse reference noted in this paper under the other Capuse, 1973a, reference and indicate it as 1973b.
- p. 75. Add the following reference under Meyrick and change the listed 1937 reference to 1937a: 1937b. Aegeriadae. *Exotic Microlepidoptera* 5:119. Add the Minet reference as listed below under literature cited.

#### ACKNOWLEDGMENT

We thank P. Leraut, Saint-Maur, France, for noting some of the errors corrected above.

#### LITERATURE CITED

- CAPUSE, I. 1973. Zur Systematik und Morphologie der Typen der Sesiidae (Lepidoptera) in der R. Püngeler-Sammlung des Zoologischen Museums zu Berlin. Mitt. Münchner Ent. Ges. 63:134–171.
- HEPPNER, J. B. & W. D. DUCKWORTH. 1981. Classification of the superfamily Sesioidea (Lepidoptera: Ditrysia). Smithsonian Contr. Zool. 314:1-144.
- MINET, J. 1976. Contribution à l'étude des microlépidoptères de Madagascar. Bull. Soc. Ent. France 81:40–43.